



HEAVY METALS URINALYSIS



Urine Toxic Metals

Urine elements are traditionally used to evaluate exposure to potentially toxic elements and wasting of nutrient elements. Additionally, the comparison of urine element concentrations before and after administration of a chelator can be used to estimate net retention of potentially toxic elements. Subsequent urine element analyses, also following the administration of a chelator, are useful for monitoring the efficacy of metal detoxification therapy. Results are expressed per 24 hours or creatinine corrected to account for urine dilution effects.



Detailed Information

Analysis of the levels of toxic metals in urine after the administration of a metal detoxification agent is an objective way to evaluate the accumulation of toxic metals. Acute metal poisoning is rare. More common, however, is a chronic, low-level exposure to toxic metals that can result in significant retention in the body that can be associated with a vast array of adverse health effects and chronic disease.



THIS TEST IS USEFUL FOR

Alopecia

Bone density

Cardiovascular disease

Depression

Dermatitis or poor wound healing

Detoxification therapy

Fatigue

Gastrointestinal symptoms

Hypertension

Immune function

Impaired glucose tolerance

Inflammation

Kidney function

Nutritional deficiencies

Parkinson's-like symptoms

Toxic element exposure





One cannot draw valid conclusions about adverse health effects of metals without assessing net retention. For an individual, toxicity occurs when net retention exceeds physiological tolerance. Net retention is determined by the difference between the rates of assimilation and excretion of metals. To evaluate net retention, one compares the levels of metals in urine before and after the administration of a pharmaceutical metal detoxification agent such as EDTA, DMSA or DMPS.

Different compounds have different affinities for specific metals, but all function by sequestering “hidden” metals from deep tissue stores and mobilizing the metals to the kidneys for excretion in the urine.

It is important to perform both pre- and postprovocation urinalysis to permit distinction between ongoing exposures to metals (pre-) and net bodily retention. The pre-provocation urine collection can also be utilized to assess the rate of creatinine clearance if a serum specimen is also submitted.

Many clinicians also request the analysis of essential elements in urine specimens to evaluate nutritional status and the efficacy of mineral supplementation during metal detoxification therapy. Metal detoxification agents can significantly increase the excretion of specific nutrient elements such as zinc, copper, manganese and molybdenum.

Chromium metabolism authorities suggest that 24-hour chromium excretion likely provides the best assessment of chromium status. Early indication of renal dysfunction can be gleaned from urinary wasting of essential elements such as magnesium, calcium, potassium and sodium in an unprovoked specimen.

Variability in urine volume can drastically affect the concentration of elements. To compensate for urine dilution variation, elements are expressed per unit creatinine for timed collections. For 24-hour collections, elements are reported as both units per 24 hours and units per creatinine.



Analytes Tested

Aluminum - urine

Antimony - urine

Arsenic - urine

Barium - urine

Beryllium - urine

Bismuth - urine

Cadmium - urine

Cesium - urine

Gadolinium - urine

Lead - urine

Mercury - urine

Nickel - urine

Palladium - urine

Platinum - urine

Tellurium - urine

Thallium - urine

Thorium - urine

Tin - urine

Tungsten - urine

Uranium - urine

